




**ECD Global Alliance
Patient and Family Gathering
Houston, TX
October 9, 2015**



Importance of Clinical Trials

Filip Janku

**Investigational Cancer Therapeutics
(Phase I Clinical Trials Program)
MD Anderson Cancer Center
Houston, TX**



Why Do We Need Clinical Trials

- Clinical trials answer two important questions
 - Does the new treatment work?
 - Is the new treatment safe?

How Do We Do Clinical Trials

- Clinical trials are usually carried out in “phases”
 - **Phase I:** What is the safe dose?
 - **Phase II:** Does the treatment work?
 - **Phase III:** Is the treatment better than existing options?
- Timeline: 10-15 years

Clinical Trials: Pros

- **For mankind**

- Increase in knowledge about particular disease and its therapy
- Development of new therapies
- Prove of efficacy or lack of thereof
- Identification of potential significant side effects

- **For individual patient**

- Access to new therapies, which are not commercially available
- Expansion of therapeutic options
- Standardized protocol-driven therapy
- Some studies in cancer patients suggested that patients on clinical trials tend to have better outcomes compared to patients treated outside of trials

Clinical Trials: Cons

- **For mankind**
 - None
- **For individual patient**
 - Need to meet all qualifying criteria, which are usually not flexible
 - Less flexible and often more intense schedule
 - Travel, financial consequences and time commitment
 - Possible risk of unknown/unexpected side effects

Why Should I Consider Clinical Trials

- Results of clinical trials are important not only for developing new therapies
- Clinical trials can provide necessary evidence to convince payers to reimburse new and effective therapies
- Clinical testing is necessary tool to make the progress happen

Strategies for Clinical Trials in ECD

- Prognosis and outcomes have dramatically improved; however, overall there is still room for improvement
- We have relatively limited therapeutic armamentarium
- We have limited resources (patients, finances) and large number of questions, which need to be answered

Strategies for Clinical Trials in ECD

- **Phase I:** Access for ECD patients to these studies, which are often limited to conventional cancers
- **Phase II:**
 - “Basket studies”: clinical trials for patients with any cancer or histiocytosis with certain unifying feature (e.g. vemurafenib in patients with *BRAF* mutation)
 - ECD specific phase II studies: because of limited number of patients this approach should be reserved for promising therapies with high likelihood of FDA/EMA approval
- **Phase III:** not feasible in ECD

Where Can I Learn About Clinical Trials?

- ECD Global Alliance Website
- [Clinicaltrials.gov](https://clinicaltrials.gov)
- Care Centers

Examples of Clinical Trials for ECD Patients

- *BRAF* mutation positive
 - **NIH: dabrafenib/trametinib**
 - Multicenter: my pathway (vemurafenib)
 - Multicenter: dabrafenib/trametinib
 - Multicenter: BVD-523
 - Multicenter: PLX8394
- *BRAF* mutation negative
 - Multicenter: BVD-523 (if MAP2K1 mutation present)
 - Multicenter: PLX8394
 - Single Center: everolimus/anakinra

Take Home Message

Clinical Trials are Part of Standard of Care in ECD